

## **Detecting single gravitons with quantum sensing**

Germain Tobar<sup>1</sup>, Sreenath K Manikandan<sup>1,2</sup>, Thomas Beitel<sup>3</sup>, and Igor Pikovski<sup>1,3</sup>

<sup>1</sup>*Stockholm University, Roslagstullsbacken 21, Stockholm, Sweden*

<sup>2</sup>*Nordita, KTH Royal Institute of Technology and Stockholm University*

<sup>3</sup>*Department of Physics, Stevens Institute of Technology, Hoboken, New Jersey 07030, USA*

The quantization of gravity is widely believed to result in gravitons - particles of discrete energy that form gravitational waves. But their detection has so far been considered impossible. Here we show that signatures of single gravitons can be observed in laboratory experiments. We show that stimulated and spontaneous singlegraviton processes can become relevant for massive quantum acoustic resonators and that stimulated absorption can be resolved through continuous sensing of quantum jumps. We analyze the feasibility of observing the exchange of single energy quanta between matter and gravitational waves. Our results show that single graviton signatures are within reach of experiments. In analogy to the discovery of the photoelectric effect for photons, such signatures can provide the first experimental evidence of the quantization of gravity.

- [1] F. Dyson, *International journal of modern physics. A, Particles and fields, gravitation, cosmology* 28, 1330041 (2013)
- [2] T. Rothman and S. Boughn, *Foundations of Physics* 36, 1801 (2006)
- [3] S. Boughn and T. Rothman, *Classical and quantum gravity* 23, 5839 (2006)
- [4] L. P. Grishchuk, *Phys. Rev. D* 45, 2601 (1992)
- [5] W. E. Lamb Jr and M. O. Scully, in *Polarization, Matter, Radiation* (Presses Universitaires, Paris, 1968) pp. 363–369